SUPPLEMENTARY TABLE 3

Bilateral-presentation cohort: descriptive profile and intraoperative findings (n = 52)

Panel A. Clinical profile and perioperative course				
Characteristic	n	%	Median	IQR
Age (month)			48	12-72
Time from presentation to surgery (days)			10	4-30.5
Follow-up (month)			24	24-27.2
Initial bilateral exploration	52	100.0		
Bilateral intraoperative confirmation of inguinal hernia	52	100.0		
Any sliding-positive hernia content	13	25.0		
Bilateral sliding-positive hernia content	3	5.8		
Second groin operation during follow-up	0	0.0		
Panel B. Intraoperative hernia sac content at index operation by side				
	Right side		Left side	
Hernia sac content	n	%	n	%
Empty sac	46	88.5	42	80.8
Ovary	6	11.5	6	11.5
Fallopian tube	0	0.0	1	1.9
Ovary + fallopian tube	0	0.0	0	0.0
Omentum	0	0.0	3	5.8

IQR, interquartile range. This cohort was excluded from the primary comparative analysis by design and is reported descriptively. Sliding-positive content denotes ovary, fallopian tube, ovary plus fallopian tube, or omentum. Panel B percentages use n = 52 per side. No missing data were present for the variables shown.